

## Exponential and Logarithmic Equations

Properties of Logarithms with base  $b$ , where  $b, x, y > 0$ ,  $b \neq 1$ , and  $p$  is a real number

1.  $\log_b 1 = 0$

$b^0 = 1$

2.  $\log_b b = 1$

$b^1 = b$

3.  $\log_b b^x = x$

4.  $b^{\log_b x} = x$

5)  $\log_b(x \cdot y) = \log_b x + \log_b y$

product property

6)  $\log_b \left( \frac{x}{y} \right) = \log_b x - \log_b y$

quotient property

7)  $\log_b(x^p) = p \cdot \log_b x$

power property

## Solving Exponential Equations

- isolate the exponential expression ✓
- take the log of both sides ✓
- simplify using log properties
- solve the equation by isolating the variable ✓

$x, t, \dots$

1)  $5^{2x} = 3$

$\log 5^{2x} = \log 3$

power property

$2x \cdot \log 5 = \log 3$

$x = \frac{\log 3}{2 \log 5}$

exact answer

$x \approx 0.341$   
approx answer

# Solving Exponential Equations

✓ 1.  $5^{2x} = 3$

4.  $e^{x^2+x} = e^{3x+15}$

✓ 2.  $160e^{-0.03t} = 80$   $\div 160$

$$e^{-0.03t} = \frac{80}{160}$$

3.  $96 - 10^x = 12$

$$e^{-0.03t} = \frac{1}{2} \quad | \text{ take ln both sides}$$

$$\ln e^{-0.03t} = \ln \frac{1}{2}$$

$$-0.03t \ln e = \ln \left( \frac{1}{2} \right)$$

$$-0.03t = \ln \left( \frac{1}{2} \right) \quad | \div -0.03$$

$$t = \frac{\ln \left( \frac{1}{2} \right)}{-0.03}$$

$$t = 23.105$$

5.  $(1 + 0.05/4)^{4t} = 3$

$$5. \log \left( 1 + \frac{0.05}{4} \right)^{4t} = \log 3$$

$$4t \cdot \log \left( 1 + \frac{0.05}{4} \right) = \log 3$$

$$t = \frac{\log 3}{4 \log \left( 1 + \frac{0.05}{4} \right)}$$

③  $96 - 10^x = 12$

$$-10^x = 12 - 96$$

$$-10^x = -84$$

$$10^x = 84$$

$$\log 10^x = \log 84$$

$$x \log 10 = \log 84$$
$$x = \log 84$$

④  $e^{x^2+x} = e^{3x+15}$

$$x^2 + x = 3x + 15$$

$$x^2 - 2x - 15 = 0$$

$$(x+3)(x-5) = 0$$

$$x = -3, x = 5$$

Solving log equations (all terms have a log with the same base)

1. condense logs into a single log (first use prop 7), if multiple logs on one side
2. solve the equation using the arguments
3. check and discard extraneous solutions

**EX** Solve the logarithmic equation algebraically.

1.  $\log(2x + 4) = \log(3 - x)$

$$2x + 4 = 3 - x$$
$$3x = -1$$

$$\rightarrow x = -\frac{1}{3}$$

check:

$$\log(2(-\frac{1}{3}) + 4) \stackrel{?}{=} \log(3 - (-\frac{1}{3}))$$
$$\log(-\frac{2}{3} + 4) \stackrel{?}{=} \log(3 + \frac{1}{3})$$
$$\log \frac{10}{3} \stackrel{?}{=} \log \frac{10}{3} \checkmark$$

2.  $\ln(x) + \ln(x - 3) = \ln(5x - 7)$

$$\ln(x(x-3)) = \ln(5x-7)$$

$$x(x-3) = 5x-7$$

$$x^2 - 3x - 5x + 7 \Rightarrow x^2 - 8x + 7 = 0$$

$$(x-7)(x-1) = 0$$

$$x = 7$$

$$x = 1$$

3.  $2\log_5 x = \log_5(7) + \log_5(3x)$

$$\log_5 x^2 = \log_5(7 \cdot 3x)$$

$$x^2 = 21x$$

$$x^2 - 21x = 0$$

$$x(x-21) = 0$$

$$x = 21$$

$$x \neq 0$$

not

check

$$x = 7$$

$$\ln 7 + \ln(7-3) \stackrel{?}{=} \ln(5 \cdot 7 - 7)$$

$$\ln 7 + \ln 4 \stackrel{?}{=} \ln 28$$
$$\ln 28 = \ln 28$$

check  $x = 1$

$$\ln 1 + \ln(1-3) \stackrel{?}{=} \ln(5 \cdot 1 - 7)$$

## Solving log equations (has a term without a log)

1. isolate terms with a log of same base to one side
2. condense logs into a single log and isolate
3. write as an exponential equation
4. solve the equation
5. check and discard extraneous solutions

4.  $-3 + 4 \ln(7x) = 2$

$$4 \ln(7x) = 5$$

$$\ln(7x) = \frac{5}{4}$$

5.  $\ln(\sqrt{x-8}) = 5$

$$e^5 = \sqrt{x-8} \quad |^2$$

$$e^{10} = x-8 \Rightarrow x = e^{10} + 8$$

6.  $3 \log_2(x+4) = 7$

$$\log_2(x+4) = \frac{7}{3}$$

$$2^{\frac{7}{3}} = x+4 \Rightarrow x = 2^{\frac{7}{3}} - 4$$

7.  $\log_4(x+2) = 1 - \log_4(x-1)$

$$+\log_4(x-1) \quad +\log_4(x-1)$$

$$\log_4(x+2) + \log_4(x-1) = 1$$

$$\log_4(x+2)(x-1) = 1$$

$$4^1 = (x+2)(x-1)$$

$$x^2 - x + 2x - 2 - 4 = 0$$

$$x^2 + x - 6 = 0$$

$$(x+3)(x-2) = 0$$

$$x = -3 \quad x = 2$$

8.  $\log x = 2 + \log(x-3)$

$$\log x - \log(x-3) = 2$$

$$\log \frac{x}{x-3} = 2$$

$$10^2 = \frac{x}{x-3} \Rightarrow$$

$$100 = \frac{x}{x-3} \Rightarrow$$

$$100(x-3) = x$$

$$100x - 300 - x = 0$$

$$99x = 300 \Rightarrow x = \frac{300}{99}$$